

Package: dprop (via r-universe)

October 22, 2024

Type Package

Title Computation of Some Important Distributional Properties

Version 0.1.0

Author Christophe Chesneau [aut], Muhammad Imran [aut, cre], M.H Tahir [aut], Farrukh Jamal [aut]

Maintainer Muhammad Imran <imranshako0r84@yahoo.com>

Depends R (>= 4.0)

Imports extraDistr, stats, VaRES

Description Generally, most of the packages specify the probability density function, cumulative distribution function, quantile function, and random numbers generation of the probability distributions. The present package allows to compute some important distributional properties, including the first four ordinary and central moments, Pearson's coefficient of skewness and kurtosis, the mean and variance, coefficient of variation, median, and quartile deviation at some parametric values of several well-known and extensively used probability distributions.

License GPL-2

Encoding UTF-8

RoxygenNote 7.2.3

NeedsCompilation no

Date/Publication 2023-06-30 11:00:06 UTC

Repository <https://imranshako0r.r-universe.dev>

RemoteUrl <https://github.com/cran/dprop>

RemoteRef HEAD

RemoteSha 2929c9d9001e864865b680c3106e57508983b02d

Contents

dprop-package	2
Beta distribution	3
Beta exponential distribution	4
Birnbaum-Saunders distribution	5
Burr XII distribution	6
Chen distribution	8
Chi-squared distribution	9
Exponential distribution	10
Exponential extension distribution	11
Exponentiated exponential distribution	12
Exponentiated Weibull distribution	13
F distribution	14
Frechet distribution	15
Gamma distribution	17
Gompertz distribution	18
Gumbel distribution	19
Inverse-gamma distribution	20
Kumaraswamy Burr XII distribution	21
Kumaraswamy distribution	22
Kumaraswamy exponential distribution	23
Kumaraswamy normal distribution	25
Laplace distribution	26
Log-normal distribution	27
Logistic distribution	28
Lomax distribution	29
Nakagami distribution	30
Normal distribution	31
Rayleigh distribution	32
Student's t distribution	33
Weibull distribution	35
Index	36

dprop-package

Computation of Some Important Distributional Properties

Description

Generally, most of the packages specify the probability density function, cumulative distribution function, quantile function, and random numbers generation of the probability distributions. The present package allows to compute some important distributional properties, including the first four ordinary and central moments, Pearson's coefficient of skewness and kurtosis, the mean and variance, coefficient of variation, median, and quartile deviation at some parametric values of several well-known and extensively used probability distributions.

Details

Package: dprop
Type: Package
Version: 0.1.0
Date: 2023-06-28
License: GPL-2

Maintainers

Muhammad Imran <imranshako0r84@yahoo.com>

Author(s)

Christophe Chesneau <christophe.chesneau@unicaen.fr>, Muhammad Imran <imranshako0r84@yahoo.com>, M.H Tahir <mht@iub.edu.pk> and Farrukh Jamal <farrukh.jamal@iub.edu.pk>.

Beta distribution *Compute the distributional properties of the beta distribution*

Description

Compute the first four ordinary moments, central moments, mean, variance, Pearson's coefficient of skewness, kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the beta distribution.

Usage

d_beta(alpha, beta)

Arguments

alpha The strictly positive shape parameter of the beta distribution ($\alpha > 0$).
beta The strictly positive shape parameter of the beta distribution ($\beta > 0$).

Details

The following is the probability density function of the beta distribution:

$$f(x) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1} (1-x)^{\beta-1},$$

where $0 \leq x \leq 1$, $\alpha > 0$ and $\beta > 0$.

Value

d_beta gives the first four ordinary moments, central moments, mean, variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the beta distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshako0r84@yahoo.com>.

References

Gupta, A. K., & Nadarajah, S. (2004). Handbook of beta distribution and its applications. CRC Press.

Johnson, N. L., Kotz, S., & Balakrishnan, N. (1994). Beta distributions. Continuous univariate distributions. 2nd ed. New York, NY: John Wiley and Sons, 221-235.

See Also

[d_kum](#)

Examples

```
d_beta(2,2)
```

Beta exponential distribution

Compute the distributional properties of the beta exponential distribution

Description

Compute the first four ordinary moments, central moments, mean, variance, Pearson's coefficient of skewness, kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the beta exponential distribution.

Usage

```
d_bexp(lambda, alpha, beta)
```

Arguments

lambda	The strictly positive scale parameter of the exponential distribution ($\lambda > 0$).
alpha	The strictly positive shape parameter of the beta distribution ($\alpha > 0$).
beta	The strictly positive shape parameter of the beta distribution ($\beta > 0$).

Details

The following is the probability density function of the beta exponential distribution:

$$f(x) = \frac{\lambda e^{-\beta \lambda x}}{B(\alpha, \beta)} (1 - e^{-\lambda x})^{\alpha-1},$$

where $x > 0$, $\alpha > 0$, $\beta > 0$ and $\lambda > 0$.

Value

`d_bexp` gives the first four ordinary moments, central moments, mean, variance, Pearson's coefficient of skewness, kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the beta exponential distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshako0r84@yahoo.com>.

References

Nadarajah, S., & Kotz, S. (2006). The beta exponential distribution. *Reliability Engineering & System Safety*, 91(6), 689-697.

See Also

[d_beta](#)

Examples

```
d_bexp(1, 1, 0.2)
```

Birnbau-Saunders distribution

Compute the distributional properties of the Birnbau-Saunders distribution

Description

Compute the first four ordinary moments, central moments, mean, variance, Pearson's coefficient of skewness, kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Birnbau-Saunders distribution.

Usage

```
d_bs(v)
```

Arguments

`v` The strictly positive scale parameter of the Birnbaum-Saunders distribution ($v > 0$).

Details

The following is the probability density function of the Birnbaum-Saunders distribution:

$$f(x) = \frac{x^{0.5} + x^{-0.5}}{2vx} \phi\left(\frac{x^{0.5} - x^{-0.5}}{v}\right),$$

where $x > 0$ and $v > 0$.

Value

`d_bs` gives the first four ordinary moments, central moments, mean, variance, Pearson's coefficient of skewness, kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Birnbaum-Saunders distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshako0r84@yahoo.com>.

References

Chan, S., Nadarajah, S., & Afuecheta, E. (2016). An R package for value at risk and expected shortfall. *Communications in Statistics Simulation and Computation*, 45(9), 3416-3434.

See Also

[d_normal](#)

Examples

```
d_bs(5)
```

Burr XII distribution *Compute the distributional properties of the Burr XII distribution*

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Burr XII distribution.

Usage`d_burr(k, c)`**Arguments**

`k` The strictly positive shape parameter of the Burr XII distribution ($k > 0$).
`c` The strictly positive shape parameter of the Burr XII distribution ($c > 0$).

Details

The following is the probability density function of the Burr XII distribution:

$$f(x) = kcx^{c-1} (1 + x^c)^{-k-1},$$

where $x > 0$, $c > 0$ and $k > 0$.

Value

`d_burr` gives the first four ordinary moments, central moments, mean, variance, Pearson's coefficient of skewness, kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Burr XII distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshakoor84@yahoo.com>.

References

Rodriguez, R. N. (1977). A guide to the Burr type XII distributions. *Biometrika*, 64(1), 129-134.
Zimmer, W. J., Keats, J. B., & Wang, F. K. (1998). The Burr XII distribution in reliability analysis. *Journal of Quality Technology*, 30(4), 386-394.

See Also

[d_kburr](#)

Examples

```
d_burr(2,10)
```

Chen distribution *Compute the distributional properties of the Chen distribution*

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Chen distribution.

Usage

`d_chen(k, c)`

Arguments

`k` The strictly positive shape parameter of the Chen distribution ($k > 0$).
`c` The strictly positive scale parameter of the Chen distribution ($c > 0$).

Details

The following is the probability density function of the Chen distribution:

$$f(x) = ckx^{k-1}e^{x^k}e^{c-ce^{x^k}},$$

where $x > 0$, $c > 0$ and $k > 0$.

Value

`d_chen` gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Chen distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshakoor84@yahoo.com>.

References

Chen, Z. (2000). A new two-parameter lifetime distribution with bathtub shape or increasing failure rate function. *Statistics & Probability Letters*, 49(2), 155–161.

See Also

[d_wei](#), [d_EE](#), [d_EW](#)

Examples

```
d_chen(0.2, 0.2)
```

Chi-squared distribution

Compute the distributional properties of the Chi-squared distribution

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the (non-central) Chi-squared distribution.

Usage

```
d_chi(n)
```

Arguments

`n` It is a degree of freedom and the positive parameter of the Chi-squared distribution ($n > 0$).

Details

The following is the probability density function of the (non-central) Chi-squared distribution:

$$f(x) = \frac{1}{2^{\frac{n}{2}} \Gamma(\frac{n}{2})} x^{\frac{n}{2}-1} e^{-\frac{x}{2}},$$

where $x > 0$ and $n > 0$.

Value

`d_chi` gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the (non-central) Chi-squared distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshakoar84@yahoo.com>.

References

- Ding, C. G. (1992). Algorithm AS275: computing the non-central chi-squared distribution function. *Journal of the Royal Statistical Society. Series C (Applied Statistics)*, 41(2), 478-482.
- Johnson, N. L., Kotz, S., & Balakrishnan, N. (1995). *Continuous univariate distributions, volume 2* (Vol. 289). John Wiley & Sons.

See Also[d_gamma](#)**Examples**`d_chi(2)`

Exponential distribution

Compute the distributional properties of the exponential distribution

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the exponential distribution.

Usage`d_exp(alpha)`**Arguments**

`alpha` The strictly positive scale parameter of the exponential distribution ($\alpha > 0$).

Details

The following is the probability density function of the exponential distribution:

$$f(x) = \alpha e^{-\alpha x},$$

where $x > 0$ and $\alpha > 0$.

Value

`d_exp` gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the exponential distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshakoor84@yahoo.com>.

References

Balakrishnan, K. (2019). Exponential distribution: theory, methods and applications. Routledge.

Singh, A. K. (1997). The exponential distribution-theory, methods and applications, *Technometrics*, 39(3), 341-341.

See Also[d_wei](#), [d_EE](#)**Examples**`d_exp(2)`

Exponential extension distribution

Compute the distributional properties of the exponential extension distribution

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the exponential extension distribution.

Usage`d_nh(alpha, beta)`**Arguments**

`alpha` The strictly positive parameter of the exponential extension distribution ($\alpha > 0$).
`beta` The strictly positive parameter of the exponential extension distribution ($\beta > 0$).

Details

The following is the probability density function of the exponential extension distribution:

$$f(x) = \alpha\beta(1 + \alpha x)^{\beta-1}e^{1-(1+\alpha x)^\beta},$$

where $x > 0$, $\alpha > 0$ and $\beta > 0$.

Value

`d_nh` gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the exponential extension distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshakoor84@yahoo.com>.

References

Nadarajah, S., & Haghighi, F. (2011). An extension of the exponential distribution. *Statistics*, 45(6), 543-558.

See Also

[d_exp](#)

Examples

`d_nh(0.5, 1)`

Exponentiated exponential distribution

Compute the distributional properties of the exponentiated exponential distribution

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the exponentiated exponential distribution.

Usage

`d_EE(alpha, beta)`

Arguments

alpha	The strictly positive scale parameter of the exponential distribution ($\alpha > 0$).
beta	The strictly positive shape parameter of the exponentiated exponential distribution ($\beta > 0$).

Details

The following is the probability density function of the exponentiated exponential distribution:

$$f(x) = \alpha\beta e^{-\alpha x} (1 - e^{-\alpha x})^{\beta-1},$$

where $x > 0$, $\alpha > 0$ and $\beta > 0$.

Value

`d_EE` gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the exponentiated exponential distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshako0r84@yahoo.com>.

References

Nadarajah, S. (2011). The exponentiated exponential distribution: a survey. *AStA Advances in Statistical Analysis*, 95, 219-251.

Gupta, R. D., & Kundu, D. (2007). Generalized exponential distribution: Existing results and some recent developments. *Journal of Statistical Planning and Inference*, 137(11), 3537-3547.

See Also

[d_EW](#), [d_wei](#), [d_exp](#)

Examples

d_EE(5,2)

Exponentiated Weibull distribution

Compute the distributional properties of the exponentiated Weibull distribution

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the exponentiated Weibull distribution.

Usage

d_EW(a, beta, zeta)

Arguments

a	The strictly positive shape parameter of the exponentiated Weibull distribution ($a > 0$).
beta	The strictly positive scale parameter of the baseline Weibull distribution ($\beta > 0$).
zeta	The strictly positive shape parameter of the baseline Weibull distribution ($\zeta > 0$).

Details

The following is the probability density function of the exponentiated Weibull distribution:

$$f(x) = a\zeta\beta^{-\zeta}x^{\zeta-1}e^{-\left(\frac{x}{\beta}\right)^\zeta} \left[1 - e^{-\left(\frac{x}{\beta}\right)^\zeta}\right]^{a-1},$$

where $x > 0$, $a > 0$, $\beta > 0$ and $\zeta > 0$.

Value

d_EW gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the exponentiated Weibull distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshakoor84@yahoo.com>.

References

Nadarajah, S., Cordeiro, G. M., & Ortega, E. M. (2013). The exponentiated Weibull distribution: a survey. *Statistical Papers*, 54, 839-877.

See Also

[d_EE](#), [d_wei](#)

Examples

d_EW(1, 1, 0.5)

F distribution

Compute the distributional properties of the F distribution

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the F distribution.

Usage

d_F(alpha, beta)

Arguments

alpha	The strictly positive parameter of the F distribution ($\alpha > 0$).
beta	The strictly positive parameter of the F distribution ($\beta > 0$).

Details

The following is the probability density function of the F distribution:

$$f(x) = \frac{1}{B\left(\frac{\alpha}{2}, \frac{\beta}{2}\right)} \left(\frac{\alpha}{\beta}\right)^{\frac{\alpha}{2}} x^{\frac{\alpha}{2}-1} \left(1 + \frac{\alpha}{\beta}x\right)^{-\left(\frac{\alpha+\beta}{2}\right)},$$

where $x > 0$, $\alpha > 0$ and $\beta > 0$.

Value

d_F gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the F distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshako0r84@yahoo.com>.

References

Johnson, N. L., Kotz, S., & Balakrishnan, N. (1995). Continuous univariate distributions, volume 2 (Vol. 289). John Wiley & Sons.

See Also

[d_gamma](#)

Examples

d_F(2,10)

Frechet distribution *Compute the distributional properties of the Frechet distribution*

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Frechet distribution.

Usage

```
d_fre(alpha, beta, zeta)
```

Arguments

alpha	The parameter of the Frechet distribution ($\alpha > 0$).
beta	The parameter of the Frechet distribution ($\beta \in (-\infty, +\infty)$).
zeta	The parameter of the Frechet distribution ($\zeta > 0$).

Details

The following is the probability density function of the Frechet distribution:

$$f(x) = \frac{\alpha}{\zeta} \left(\frac{x - \beta}{\zeta} \right)^{-1-\alpha} e^{-\left(\frac{x-\beta}{\zeta}\right)^{-\alpha}},$$

where $x > \beta$, $\alpha > 0$, $\zeta > 0$ and $\beta \in (-\infty, +\infty)$. The Frechet distribution is also known as inverse Weibull distribution and special case of the generalized extreme value distribution.

Value

d_fre gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Frechet distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshako0r84@yahoo.com>.

References

Abbas, K., & Tang, Y. (2015). Analysis of Frechet distribution using reference priors. *Communications in Statistics-Theory and Methods*, 44(14), 2945-2956.

See Also

[d_wei](#)

Examples

```
d_fre(5, 1, 0.5)
```

Gamma distribution *Compute the distributional properties of the gamma distribution*

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the gamma distribution.

Usage

```
d_gamma(alpha, beta)
```

Arguments

alpha The strictly positive parameter of the gamma distribution ($\alpha > 0$).
beta The strictly positive parameter of the gamma distribution ($\beta > 0$).

Details

The following is the probability density function of the gamma distribution:

$$f(x) = \frac{\beta^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\beta x},$$

where $x > 0$, $\alpha > 0$ and $\beta > 0$.

Value

d_gamma the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the gamma distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshakoor84@yahoo.com>.

References

Burgin, T. A. (1975). The gamma distribution and inventory control. Journal of the Operational Research Society, 26(3), 507-525.

See Also

[d_wei](#), [d_naka](#)

Examples

```
d_gamma(2, 2)
```

Gompertz distribution *Compute the distributional properties of the Gompertz distribution*

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Gompertz distribution.

Usage

```
d_gompertz(alpha, beta)
```

Arguments

alpha	The strictly positive parameter of the Gompertz distribution ($\alpha > 0$).
beta	The strictly positive parameter of the Gompertz distribution ($\beta > 0$).

Details

The following is the probability density function of the Gompertz distribution:

$$f(x) = \alpha e^{\beta x - \frac{\alpha}{\beta}(e^{\beta x} - 1)},$$

where $x > 0$, $\alpha > 0$ and $\beta > 0$.

Value

`d_gompertz` gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Gompertz distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshakoor84@yahoo.com>.

References

Soliman, A. A., Abd-Allah, A. H., Abou-Elheggag, N. A., & Abd-Elmougod, G. A. (2012). Estimation of the parameters of life for Gompertz distribution using progressive first-failure censored data. *Computational Statistics & Data Analysis*, 56(8), 2471-2485.

See Also

[d_fre](#)

Examples

```
d_gompertz(2,2)
```

Gumbel distribution *Compute the distributional properties of the Gumbel distribution*

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Gumbel distribution.

Usage

```
d_gumbel(alpha, beta)
```

Arguments

alpha	Location parameter of the Gumbel distribution ($\alpha \in (-\infty, +\infty)$).
beta	The strictly positive scale parameter of the Gumbel distribution ($\beta > 0$).

Details

The following is the probability density function of the Gumbel distribution:

$$f(x) = \frac{1}{\beta} e^{-(z+e^{-z})},$$

where $z = \frac{x-\alpha}{\beta}$, $x \in (-\infty, +\infty)$, $\alpha \in (-\infty, +\infty)$ and $\beta > 0$.

Value

`d_gumbel` gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Gumbel distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshakoor84@yahoo.com>.

References

Gomez, Y. M., Bolfarine, H., & Gomez, H. W. (2019). Gumbel distribution with heavy tails and applications to environmental data. *Mathematics and Computers in Simulation*, 157, 115-129.

See Also

[d_gompertz](#), [d_fre](#)

Examples

```
d_gumbel(1,2)
```

Inverse-gamma distribution

Compute the distributional properties of the inverse-gamma distribution

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the inverse-gamma distribution.

Usage

```
d_ingam(alpha, beta)
```

Arguments

alpha	The strictly positive parameter of the inverse-gamma distribution ($\alpha > 0$).
beta	The strictly positive parameter of the inverse-gamma distribution ($\beta > 0$).

Details

The following is the probability density function of the inverse-gamma distribution:

$$f(x) = \frac{\beta^\alpha}{\Gamma(\alpha)} x^{-\alpha-1} e^{-\frac{\beta}{x}},$$

where $x > 0$, $\alpha > 0$ and $\beta > 0$.

Value

`d_ingam` gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the inverse-gamma distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshakoor84@yahoo.com>.

References

- Rivera, P. A., Calderín-Ojeda, E., Gallardo, D. I., & Gómez, H. W. (2021). A compound class of the inverse Gamma and power series distributions. *Symmetry*, 13(8), 1328.
- Glen, A. G. (2017). On the inverse gamma as a survival distribution. *Computational Probability Applications*, 15-30.

See Also

[d_gamma](#)

Examples

`d_ingam(5,2)`

Kumaraswamy Burr XII distribution

Compute the distributional properties of the Kumaraswamy Burr XII distribution

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Kumaraswamy Burr XII distribution.

Usage

`d_kburr(a, b, k, c)`

Arguments

- | | |
|---|--|
| a | The strictly positive parameter of the Kumaraswamy distribution ($a > 0$). |
| b | The strictly positive parameter of the Kumaraswamy distribution ($b > 0$). |
| k | The strictly positive parameter of the Burr XII distribution ($k > 0$). |
| c | The strictly positive parameter of the Burr XII distribution ($c > 0$). |

Details

The following is the probability density function of the Kumaraswamy Burr XII distribution:

$$f(x) = \frac{abkcx^{c-1}}{(1+x^c)^{k+1}} \left[1 - (1+x^c)^{-k}\right]^{a-1} \left\{1 - \left[1 - (1+x^c)^{-k}\right]^a\right\}^{b-1},$$

where $x > 0$, $a > 0$, $b > 0$, $k > 0$ and $c > 0$.

Value

`d_kburr` gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Kumaraswamy Burr XII distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshako0r84@yahoo.com>.

References

Paranaiba, P. F., Ortega, E. M., Cordeiro, G. M., & Pascoa, M. A. D. (2013). The Kumaraswamy Burr XII distribution: theory and practice. *Journal of Statistical Computation and Simulation*, 83(11), 2117-2143.

See Also

[d_kum](#), [d_kexp](#)

Examples

```
d_kburr(1.5, 1, 1, 7)
```

Kumaraswamy distribution

Compute the distributional properties of the Kumaraswamy distribution

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Kumaraswamy distribution.

Usage

```
d_kum(alpha, beta)
```

Arguments

<code>alpha</code>	The strictly positive parameter of the Kumaraswamy distribution ($\alpha > 0$).
<code>beta</code>	The strictly positive parameter of the Kumaraswamy distribution ($\beta > 0$).

Details

The following is the probability density function of the Kumaraswamy distribution:

$$f(x) = \alpha\beta x^{\alpha-1} (1 - x^\alpha)^{\beta-1},$$

where $0 \leq x \leq 1$, $\alpha > 0$ and $\beta > 0$.

Value

`d_kum` gives the first four ordinary moments, central moments, mean, variance, Pearson's coefficient of skewness, kurtosis, coefficient of variation, median and quartile deviation at some parametric values based on the Kumaraswamy distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshakoor84@yahoo.com>.

References

El-Sherpieny, E. S. A., & Ahmed, M. A. (2014). On the kumaraswamy distribution. *International Journal of Basic and Applied Sciences*, 3(4), 372.

Mitnik, P. A. (2013). New properties of the Kumaraswamy distribution. *Communications in Statistics-Theory and Methods*, 42(5), 741-755.

Dey, S., Mazucheli, J., & Nadarajah, S. (2018). Kumaraswamy distribution: different methods of estimation. *Computational and Applied Mathematics*, 37, 2094-2111.

See Also

[d_beta](#)

Examples

```
d_kum(2, 2)
```

Kumaraswamy exponential distribution

Compute the distributional properties of the Kumaraswamy exponential distribution

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Kumaraswamy exponential distribution.

Usage

```
d_kexp(lambda, a, b)
```

Arguments

a The strictly positive shape parameter of the Kumaraswamy distribution ($a > 0$).

b The strictly positive shape parameter of the Kumaraswamy distribution ($b > 0$).

lambda The strictly positive parameter of the exponential distribution ($\lambda > 0$).

Details

The following is the probability density function of the Kumaraswamy exponential distribution:

$$f(x) = ab\lambda e^{-\lambda x} (1 - e^{-\lambda x})^{a-1} \left\{ 1 - (1 - e^{-\lambda x})^a \right\}^{b-1},$$

where $x > 0$, $a > 0$, $b > 0$ and $\lambda > 0$.

Value

`d_kexp` gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Kumaraswamy exponential distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshako0r84@yahoo.com>.

References

Cordeiro, G. M., & de Castro, M. (2011). A new family of generalized distributions. *Journal of Statistical Computation and Simulation*, 81(7), 883-898.

See Also

[d_kburr](#), [d_kum](#)

Examples

```
d_kexp(0.2, 1, 1)
```

Kumaraswamy normal distribution

Compute the distributional properties of the Kumaraswamy normal distribution

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Kumaraswamy normal distribution.

Usage

`d_kumnorm(mu, sigma, a, b)`

Arguments

<code>mu</code>	The location parameter of the normal distribution ($\mu \in (-\infty, +\infty)$).
<code>sigma</code>	The strictly positive scale parameter of the normal distribution ($\sigma > 0$).
<code>a</code>	The strictly positive shape parameter of the Kumaraswamy distribution ($a > 0$).
<code>b</code>	The strictly positive shape parameter of the Kumaraswamy distribution ($b > 0$).

Details

The following is the probability density function of the Kumaraswamy normal distribution:

$$f(x) = \frac{ab}{\sigma} \phi\left(\frac{x-\mu}{\sigma}\right) \left[\Phi\left(\frac{x-\mu}{\sigma}\right)\right]^{a-1} \left[1 - \Phi\left(\frac{x-\mu}{\sigma}\right)\right]^{b-1},$$

where $x \in (-\infty, +\infty)$, $\mu \in (-\infty, +\infty)$, $\sigma > 0$, $a > 0$ and $b > 0$. The functions $\phi(\cdot)$ and $\Phi(\cdot)$, denote the probability density function and cumulative distribution function of the standard normal variable, respectively.

Value

`d_kumnorm` gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Kumaraswamy normal distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshako0r84@yahoo.com>.

References

Cordeiro, G. M., & de Castro, M. (2011). A new family of generalized distributions. *Journal of Statistical Computation and Simulation*, 81(7), 883-898.

See Also

[d_kburr](#), [d_kexp](#), [d_kum](#)

Examples

```
d_kumnorm(0.2, 0.2, 2, 2)
```

Laplace distribution *Compute the distributional properties of the Laplace or double exponential distribution*

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Laplace distribution.

Usage

```
d_lap(alpha, beta)
```

Arguments

alpha Location parameter of the Laplace distribution ($\alpha \in (-\infty, +\infty)$).

beta The strictly positive scale parameter of the Laplace distribution ($\beta > 0$).

Details

The following is the probability density function of the Laplace distribution:

$$f(x) = \frac{1}{2\beta} e^{-\frac{|x-\alpha|}{\beta}},$$

where $x \in (-\infty, +\infty)$, $\alpha \in (-\infty, +\infty)$ and $\beta > 0$.

Value

`d_lap` gives the first four ordinary moments, central moments, mean, variance, Pearson's coefficient of skewness, kurtosis, coefficient of variation, median and quartile deviation at some parametric values based on the Laplace distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshakoor84@yahoo.com>.

References

Cordeiro, G. M., & Lemonte, A. J. (2011). The beta Laplace distribution. *Statistics & Probability Letters*, 81(8), 973-982.

See Also

[d_normal](#)

Examples

d_lap(2,4)

Log-normal distribution

Compute the distributional properties of the log-normal distribution

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the log-normal distribution.

Usage

d_lnormal(mu, sigma)

Arguments

mu The location parameter ($\mu \in (-\infty, +\infty)$).

sigma The strictly positive scale parameter of the log-normal distribution ($\sigma > 0$).

Details

The following is the probability density function of the log-normal distribution:

$$f(x) = \frac{1}{x\sigma\sqrt{2\pi}} e^{-\frac{(\log(x)-\mu)^2}{2\sigma^2}},$$

where $x > 0$, $\mu \in (-\infty, +\infty)$ and $\sigma > 0$.

Value

d_lnormal gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the log-normal distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshakoor84@yahoo.com>.

References

Johnson, N. L., Kotz, S., & Balakrishnan, N. (1995). Continuous Univariate Distributions, Volume 1, Chapter 14. Wiley, New York.

See Also

[d_normal](#)

Examples

```
d_lnormal(1,0.5)
```

Logistic distribution *Compute the distributional properties of the logistic distribution*

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the logistic distribution.

Usage

```
d_logis(mu, sigma)
```

Arguments

mu	Location parameter of the logistic distribution ($\mu \in (-\infty, +\infty)$).
sigma	The strictly positive scale parameter of the logistic distribution ($\sigma > 0$).

Details

The following is the probability density function of the logistic distribution:

$$f(x) = \frac{e^{-\frac{(x-\mu)}{\sigma}}}{\sigma \left(1 + e^{-\frac{(x-\mu)}{\sigma}}\right)^2},$$

where $x \in (-\infty, +\infty)$, $\mu \in (-\infty, +\infty)$ and $\sigma > 0$.

Value

`d_logis` gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the logistic distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshakoor84@yahoo.com>.

References

Johnson, N. L., Kotz, S., & Balakrishnan, N. (1995). Continuous univariate distributions, Volume 2 (Vol. 289). John Wiley & Sons.

See Also

[d_lnormal](#)

Examples

```
d_logis(4,0.2)
```

Lomax distribution *Compute the distributional properties of the Lomax distribution*

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Lomax distribution.

Usage

```
d_lom(alpha, beta)
```

Arguments

alpha The strictly positive parameter of the Lomax distribution ($\alpha > 0$).

beta The strictly positive parameter of the Lomax distribution ($\beta > 0$).

Details

The following is the probability density function of the Lomax distribution:

$$f(x) = \frac{\alpha}{\beta} \left(1 + \frac{x}{\beta}\right)^{-\alpha-1},$$

where $x > 0$, $\alpha > 0$ and $\beta > 0$.

Value

d_lom gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Lomax distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshakoor84@yahoo.com>.

References

Abd-Elfattah, A. M., Alaboud, F. M., & Alharby, A. H. (2007). On sample size estimation for Lomax distribution. Australian Journal of Basic and Applied Sciences, 1(4), 373-378.

See Also

[d_gamma](#)

Examples

```
d_lom(10,10)
```

Nakagami distribution *Compute the distributional properties of the Nakagami distribution*

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Nakagami distribution.

Usage

```
d_naka(alpha, beta)
```

Arguments

alpha	The strictly positive parameter of the Nakagami distribution ($\alpha > 0$).
beta	The strictly positive parameter of the Nakagami distribution ($\beta > 0$).

Details

The following is the probability density function of the Nakagami distribution:

$$f(x) = \frac{2\alpha^\alpha}{\Gamma(\alpha)\beta^\alpha} x^{2\alpha-1} e^{-\frac{\alpha x^2}{\beta}},$$

where $x > 0$, $\alpha > 0$ and $\beta > 0$.

Value

`d_naka` gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Nakagami distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshako0r84@yahoo.com>.

References

Schwartz, J., Godwin, R. T., & Giles, D. E. (2013). Improved maximum-likelihood estimation of the shape parameter in the Nakagami distribution. *Journal of Statistical Computation and Simulation*, 83(3), 434-445.

See Also

[d_gamma](#)

Examples

```
d_naka(2,2)
```

Normal distribution *Compute the distributional properties of the normal distribution*

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the normal distribution.

Usage

```
d_normal(alpha, beta)
```

Arguments

<code>alpha</code>	Location parameter of the normal distribution ($\alpha \in (-\infty, +\infty)$).
<code>beta</code>	The strictly positive scale parameter of the normal distribution ($\beta > 0$).

Details

The following is the probability density function of the normal distribution:

$$f(x) = \frac{1}{\beta\sqrt{2\pi}} e^{-0.5\left(\frac{x-\alpha}{\beta}\right)^2},$$

where $x \in (-\infty, +\infty)$, $\alpha \in (-\infty, +\infty)$ and $\beta > 0$. The parameters α and β represent the mean and standard deviation, respectively.

Value

`d_normal` gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the normal distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshakoor84@yahoo.com>.

References

Patel, J. K., & Read, C. B. (1996). Handbook of the normal distribution (Vol. 150). CRC Press.

See Also

[d_lnormal](#)

Examples

```
d_normal(4,0.2)
```

Rayleigh distribution *Compute the distributional properties of the Rayleigh distribution*

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Rayleigh distribution.

Usage

```
d_rayl(alpha)
```

Arguments

`alpha` The strictly positive parameter of the Rayleigh distribution ($\alpha > 0$).

Details

The following is the probability density function of the Rayleigh distribution:

$$f(x) = \frac{x}{\alpha^2} e^{-\frac{x^2}{2\alpha^2}},$$

where $x > 0, \alpha > 0$.

Value

`d_rayl` gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Rayleigh distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshako0r84@yahoo.com>.

References

Forbes, C., Evans, M. Hastings, N., & Peacock, B. (2011). Statistical Distributions. John Wiley & Sons.

See Also

[d_wei](#)

Examples

```
d_rayl(2)
```

Student's t distribution

Compute the distributional properties of the Student distribution

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Student t distribution.

Usage

```
d_st(v)
```

Arguments

`v` The strictly positive parameter of the Student distribution ($v > 0$), it is also called a degree of freedom.

Details

The following is the probability density function of the Student *t* distribution:

$$f(x) = \frac{\Gamma(\frac{v+1}{2})}{\sqrt{v\pi}\Gamma(\frac{v}{2})} \left(1 + \frac{x^2}{v}\right)^{-(v+1)/2},$$

where $x \in (-\infty, +\infty)$ and $v > 0$.

Value

`d_st` gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Student *t* distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshako0r84@yahoo.com>.

References

Yang, Z., Fang, K. T., & Kotz, S. (2007). On the Student's *t*-distribution and the *t*-statistic. *Journal of Multivariate Analysis*, 98(6), 1293-1304.

Ahsanullah, M., Kibria, B. G., & Shakil, M. (2014). Normal and Student's *t* distributions and their applications (Vol. 4). Paris, France: Atlantis Press.

See Also

[d_chi](#)

Examples

`d_st(6)`

Weibull distribution *Compute the distributional properties of the Weibull distribution*

Description

Compute the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Weibull distribution.

Usage

```
d_wei(alpha, beta)
```

Arguments

alpha	The strictly positive scale parameter of the Weibull distribution ($\alpha > 0$).
beta	The strictly positive shape parameter of the Weibull distribution ($\beta > 0$).

Details

The following is the probability density function of the Weibull distribution:

$$f(x) = \frac{\beta}{\alpha} \left(\frac{x}{\alpha}\right)^{\beta-1} e^{-\left(\frac{x}{\alpha}\right)^\beta},$$

where $x > 0$, $\alpha > 0$ and $\beta > 0$.

Value

d_wei gives the first four ordinary moments, central moments, mean, and variance, Pearson's coefficient of skewness and kurtosis, coefficient of variation, median and quartile deviation based on the selected parametric values of the Weibull distribution.

Author(s)

Muhammad Imran.

R implementation and documentation: Muhammad Imran <imranshako0r84@yahoo.com>.

References

Hallinan Jr, Arthur J. (1993). A review of the Weibull distribution. *Journal of Quality Technology*, 25(2), 85-93.

See Also

[d_EE](#)

Examples

```
d_wei(2,2)
```

Index

- * **Central moments**
 - dprop-package, 2
 - * **Coefficient of variation**
 - dprop-package, 2
 - * **Distribution theory**
 - dprop-package, 2
 - * **Kurtosis**
 - dprop-package, 2
 - * **Mean and variance**
 - dprop-package, 2
 - * **Median**
 - dprop-package, 2
 - * **Ordinary moments**
 - dprop-package, 2
 - * **Quartile deviation**
 - dprop-package, 2
 - * **Skewness**
 - dprop-package, 2
- Beta distribution, 3
Beta exponential distribution, 4
Birnbau-Saunders distribution, 5
Burr XII distribution, 6
- Chen distribution, 8
Chi-squared distribution, 9
- d_beta, 5, 23
d_beta (Beta distribution), 3
d_bexp (Beta exponential distribution), 4
d_bs (Birnbau-Saunders distribution), 5
d_burr (Burr XII distribution), 6
d_chen (Chen distribution), 8
d_chi, 34
d_chi (Chi-squared distribution), 9
d_EE, 8, 11, 14, 35
d_EE (Exponentiated exponential distribution), 12
d_EW, 8, 13
d_EW (Exponentiated Weibull distribution), 13
d_exp, 12, 13
d_exp (Exponential distribution), 10
d_F (F distribution), 14
d_fre, 18, 20
d_fre (Frechet distribution), 15
d_gamma, 10, 15, 21, 30, 31
d_gamma (Gamma distribution), 17
d_gompertz, 20
d_gompertz (Gompertz distribution), 18
d_gumbel (Gumbel distribution), 19
d_ingam (Inverse-gamma distribution), 20
d_kburr, 7, 24, 26
d_kburr (Kumaraswamy Burr XII distribution), 21
d_kexp, 22, 26
d_kexp (Kumaraswamy exponential distribution), 23
d_kum, 4, 22, 24, 26
d_kum (Kumaraswamy distribution), 22
d_kumnorm (Kumaraswamy normal distribution), 25
d_lap (Laplace distribution), 26
d_lnormal, 29, 32
d_lnormal (Log-normal distribution), 27
d_logis (Logistic distribution), 28
d_lom (Lomax distribution), 29
d_naka, 17
d_naka (Nakagami distribution), 30
d_nh (Exponential extension distribution), 11
d_normal, 6, 27, 28
d_normal (Normal distribution), 31
d_rayl (Rayleigh distribution), 32
d_st (Student's t distribution), 33
d_wei, 8, 11, 13, 14, 16, 17, 33
d_wei (Weibull distribution), 35
dprop-package, 2

Exponential distribution, [10](#)
Exponential extension distribution, [11](#)
Exponentiated exponential
distribution, [12](#)
Exponentiated Weibull distribution, [13](#)

F distribution, [14](#)
Frechet distribution, [15](#)

Gamma distribution, [17](#)
Gompertz distribution, [18](#)
Gumbel distribution, [19](#)

Inverse-gamma distribution, [20](#)

Kumaraswamy Burr XII distribution, [21](#)
Kumaraswamy distribution, [22](#)
Kumaraswamy exponential distribution,
[23](#)
Kumaraswamy normal distribution, [25](#)

Laplace distribution, [26](#)
Log-normal distribution, [27](#)
Logistic distribution, [28](#)
Lomax distribution, [29](#)

Nakagami distribution, [30](#)
Normal distribution, [31](#)

Rayleigh distribution, [32](#)

Student's t distribution, [33](#)

Weibull distribution, [35](#)